

A DE BRUIJN-ERDŐS THEOREM FOR 1-2 METRIC SPACES

VAŠEK CHVÁTAL, Montréal

(Received May 8, 2012)

Abstract. A special case of a combinatorial theorem of De Bruijn and Erdős asserts that every noncollinear set of n points in the plane determines at least n distinct lines. Chen and Chvátal suggested a possible generalization of this assertion in metric spaces with appropriately defined lines. We prove this generalization in all metric spaces where each nonzero distance equals 1 or 2.

Keywords: line in metric space; De Bruijn-Erdős theorem

MSC 2010: 05D99, 51G99

It is well known that

- (i) *every noncollinear set of n points in the plane determines at least n distinct lines.*

As noted by Erdős [5], theorem (i) is a corollary of the Sylvester-Gallai theorem (asserting that, for every noncollinear set S of finitely many points in the plane, some line goes through precisely two points of S); it is also a special case of a combinatorial theorem proved later by De Bruijn and Erdős [4].

Chen and Chvátal [2] suggested that theorem (i) might be generalized in the framework of metric spaces. In a Euclidean space, line \overline{uv} is characterized as

$$\overline{uv} = \{p: \text{dist}(p, u) + \text{dist}(u, v) = \text{dist}(p, v) \text{ or} \\ \text{dist}(u, p) + \text{dist}(p, v) = \text{dist}(u, v) \text{ or } \text{dist}(u, v) + \text{dist}(v, p) = \text{dist}(u, p)\},$$

where dist is the Euclidean metric; in an arbitrary metric space (S, dist) , the same relation may be taken for the definition of the line. (Unlike in the case of Euclidean

This research was undertaken, in part, thanks to funding from the Canada Research Chairs program.

lines, $x, y \in \overline{uv}$, $x \neq y$ does not imply $u, v \in \overline{xy}$; nevertheless, $x \in \overline{uv}$, $x \neq u$ still implies $v \in \overline{xu}$.) With this definition of lines in metric spaces, Chen and Chvátal asked:

- (ii) *True or false? Every metric space on n points, where $n \geq 2$, either has at least n distinct lines or else has a line that consists of all n points.*

Let us say that a metric space on n points has the *De Bruijn-Erdős property* if it either has at least n distinct lines or else has a line that consists of all n points: now we may state (ii) by asking whether or not all metric spaces on at least 2 points have the De Bruijn-Erdős property. A survey of results related to this question appears in [1].

By a 1-2 *metric space*, we mean a metric space where each nonzero distance is 1 or 2. Chiniforooshan and Chvátal [3] proved that

- (iii) *every 1-2 metric space on n points has $\Omega(n^{4/3})$ distinct lines and this bound is tight.*

This result states that all sufficiently large 1-2 metric spaces have a property far stronger than the De Bruijn-Erdős property, but it does not imply that all 1-2 metric spaces on at least 2 points have the De Bruijn-Erdős property. The purpose of the present note is to remove this blemish.

Theorem 1. *All 1-2 metric spaces on at least 2 points have the De Bruijn-Erdős property.*

The rest of this note is devoted to a proof of Theorem 1. A key notion in the proof, one borrowed from [3], is the notion of *twins* in a 1-2 metric space: these are points u, v such that $\text{dist}(u, v) = 2$ and $\text{dist}(u, w) = \text{dist}(v, w)$ for all points w distinct from both u and v . Use of this notion in counting lines is pointed out in the following claim (also borrowed from [3]), whose proof is straightforward.

Claim 1. If u_1, u_2, u_3, u_4 are four distinct points in a 1-2 metric space, then

- ▷ if $\text{dist}(u_1, u_2) \neq \text{dist}(u_3, u_4)$, then $\overline{u_1u_2} \neq \overline{u_3u_4}$,
- ▷ if $\text{dist}(u_1, u_2) = \text{dist}(u_2, u_3) = 2$, then $\overline{u_1u_2} \neq \overline{u_2u_3}$,
- ▷ if $\text{dist}(u_1, u_2) = \text{dist}(u_2, u_3) = 1$ and u_1, u_3 are not twins, then $\overline{u_1u_2} \neq \overline{u_2u_3}$.

By a *critical 1-2 metric space*, we shall mean a smallest counterexample to Theorem 1; in a sequence of claims, we shall gradually prove the nonexistence of a critical 1-2 metric space. We shall say that a line in a metric space is *universal* if, and only if, it consists of all points of the space.

Claim 2. For every pair u, v of twins in a critical 1-2 metric space, there is a third point w in this space such that $\text{dist}(u, w) = \text{dist}(v, w) = 2$ and $\text{dist}(x, y) = 1$ whenever $x \in \{u, v, w\}$, $y \notin \{u, v, w\}$.

Proof. Let S denote the space we are dealing with. Since S is critical, S does not have the De Bruijn-Erdős property and $S \setminus u$ has the De Bruijn-Erdős property. We will derive the existence of w from these two facts.

The assumption that u, v are twins implies that

(a) if x, y are distinct points in $S \setminus \{u, v\}$, then the line \overline{xy} in S contains either both u, v or neither of u, v ;

(b) if $w \in S \setminus u$ and $\text{dist}(w, v) = 1$, then the line \overline{wv} in S (and the line \overline{wu} in S) contains both u, v ;

(c) if $w \in S \setminus u$ and $\text{dist}(w, v) = 2$, then the line \overline{wv} in S contains v and not u and the line \overline{wu} in S contains u and not v .

Since S does not have the De Bruijn-Erdős property, we have $\overline{wv} \neq S$; since u and v are twins, it follows that

(d) there is a w in $S \setminus u$ such that $\text{dist}(w, v) = 2$.

From (a), (b), (c), (d), we conclude that

(e) the number of lines in S exceeds the number of lines in $S \setminus u$.

Since S does not have the De Bruijn-Erdős property, the number of lines in S is less than $|S|$, and so (e) implies that the number of lines in $S \setminus u$ is less than $|S \setminus u|$; since $S \setminus u$ has the De Bruijn-Erdős property, it follows that

(f) $S \setminus u$ has a universal line.

Since S does not have the De Bruijn-Erdős property,

(g) S has no universal line.

Facts (a), (f), and (g) together imply that some line \overline{wv} in $S \setminus u$ is universal. Now (b) and (g) together imply that $\text{dist}(w, v) = 2$; since u, v are twins, it follows that $\text{dist}(u, v) = 2$ and $\text{dist}(w, u) = 2$. Since \overline{wv} is a universal line in $S \setminus u$, we have $\text{dist}(w, y) = \text{dist}(v, y) = 1$ whenever $y \notin \{u, v, w\}$; since u, v are twins, it follows that $\text{dist}(u, y) = 1$ whenever $y \notin \{u, v, w\}$. \square

Claim 3. No critical 1-2 metric space contains a pair of twins.

Proof. Assume the contrary: some critical 1-2 metric space S contains a pair of twins. We will show that S has at least $|S|$ lines, contradicting the assumption that S does not have the De Bruijn-Erdős property. For this purpose, consider the largest set $\{T_1, T_2, \dots, T_k\}$ of pairwise disjoint three-point subsets of S such that $\text{dist}(u, v) = 2$ whenever u, v are distinct points in the same T_i and such that $\text{dist}(u, x) = 1$ whenever $u \in T_i, x \notin T_i$ for some i . Since S contains a pair of twins, Claim 2 guarantees that $k \geq 1$; we will derive the existence of $|S|$ lines in S from this fact.

Let \mathcal{L}_1 denote the set of all lines \overline{wv} such that u, v are distinct points in the same T_i . If $\overline{wv} \in \mathcal{L}_1$, then $\overline{wv} = S \setminus w$, where $\{u, v, w\} = T_i$ for some i ; it follows that

(a) \mathcal{L}_1 consists of the $3k$ sets $S \setminus w$ with w ranging through $\bigcup_{i=1}^k T_i$.

Next, choose a point r in T_1 and let \mathcal{L}_2 denote the set of all lines \overline{rx} such that $x \in S \setminus \bigcup_{i=1}^k T_i$. Claim 2 and the maximality of k together guarantee that S contains no pair x, y of twins such that $x, y \in S \setminus \bigcup_{i=1}^k T_i$. This fact and Claim 1 together imply that

(b) $|\mathcal{L}_2| = |S| - 3k$.

Finally, note that each line in \mathcal{L}_2 includes all points of T_1 and no points of T_2 . This observation and (a) together imply that $\mathcal{L}_1 \cap \mathcal{L}_2 = \emptyset$, and so $|\mathcal{L}_1 \cup \mathcal{L}_2| = |S|$ by (a) and (b). \square

Each 1-2 metric space can be thought of as a complete graph with each edge uv labeled by $\text{dist}(u, v)$. Given edges uv, xy of this complete graph, let us write $uv \approx xy$ to mean that $\overline{uv} = \overline{xy}$. The following fact is a direct consequence of Claim 1 combined with Claim 3.

Claim 4. Each equivalence class of the equivalence relation \approx in a critical 1-2 metric space is a set of pairwise disjoint edges with identical labels or else a (not necessarily proper) subset of a cycle of length four with alternating labels.

Claim 5. The size of each equivalence class of the equivalence relation \approx in a critical 1-2 metric space on n points is at most $\max\{(n-1)/2, 4\}$.

Proof. This is a direct corollary of Claim 4 combined with the observation that an equivalence class of $n/2$ pairwise disjoint edges defines a universal line. \square

Claim 6. Every critical 1-2 metric space has at most 7 points.

Proof. Consider an arbitrary critical 1-2 metric space and let n denote the number of its points. Since this space does not have the De Bruijn-Erdős property, it has fewer than n lines, and so its equivalence relation \approx partitions the $n(n-1)/2$ edges of its complete graph into at most $n-1$ classes. Since the largest of these classes has size at least $n/2$, Claim 5 implies that $n/2 \leq \max\{(n-1)/2, 4\}$, and so $n \leq 8$. If $n = 8$, then the 28 edges of the complete graph are partitioned into 7 equivalence classes of size 4. Now Claim 4 and the absence of a universal line together imply that each of these equivalence classes is a cycle of length four. But this is impossible, since the edge set of the complete graph on eight vertices cannot be partitioned into cycles: each vertex of this graph has an odd degree. \square

Claim 7. No critical 1-2 metric space has 7 points.

Proof. Consider an arbitrary critical 1-2 metric space on 7 points. Since this space does not have the De Bruijn-Erdős property, it has fewer than 7 lines, and so its equivalence relation \approx partitions the 21 edges of its complete graph into at most

6 classes. By Claim 5, each of these classes has size at most 4, and so at least three of them have size precisely 4; by Claim 4, each of these three classes is a cycle of length four. Let G_1, G_2, G_3 denote these three subgraphs of the complete graph on seven vertices.

Since G_1, G_2, G_3 are pairwise edge-disjoint, every two of them share at most two vertices; since their union has only seven vertices, some two of them share at least two vertices; we may assume (after a permutation of subscripts if necessary) that G_1 and G_2 share precisely two vertices. Let us name these two vertices u, v . Since G_1 and G_2 are edge-disjoint, we may assume (after a switch of subscripts if necessary) that vertices u, v are adjacent in G_1 and nonadjacent in G_2 .

Next, we may name w, x the remaining two vertices in G_1 in such a way that the four edges of G_1 are uv, vw, wx, ux ; we may name y, z the remaining two vertices in G_2 in such a way that the four edges of G_2 are uy, uz, vz, vy . Since the labels on the edges of G_2 alternate, we may assume (after switching y and z if necessary) that $\text{dist}(u, y) = 1, \text{dist}(u, z) = 2, \text{dist}(v, z) = 1, \text{dist}(v, y) = 2$. Since $\overline{uy} = \overline{vy}$, we have $u \in \overline{vy}$; since $\text{dist}(v, y) = 2$, it follows that $\text{dist}(u, v) = 1$. In turn, since the labels on the edges of G_1 alternate, we have $\text{dist}(v, w) = 2, \text{dist}(w, x) = 1, \text{dist}(u, x) = 2$.

Now $\text{dist}(y, u) + \text{dist}(u, v) = \text{dist}(y, v)$, and so $y \in \overline{uv}$; since $uv \approx vw$, it follows that $y \in \overline{vw}$. But this is impossible, since $\text{dist}(v, w) = 2$ and $\text{dist}(v, y) = 2$. \square

Claim 8. Every critical 1-2 metric space on 5 or 6 points contains pairwise distinct points u, v, w, x, y such that

$$\begin{aligned} \text{dist}(u, w) &= \text{dist}(u, x) = \text{dist}(v, w) = \text{dist}(v, x) = 1, \\ \text{dist}(u, v) &= \text{dist}(w, x) = 2, \\ \text{dist}(u, y) &\neq \text{dist}(v, y), \quad \text{dist}(w, y) \neq \text{dist}(x, y). \end{aligned}$$

Proof. Consider an arbitrary critical 1-2 metric space on n points such that $n = 5$ or $n = 6$. Since this space does not have the De Bruijn-Erdős property, it has fewer than n lines, and so its equivalence relation \approx partitions the $n(n-1)/2$ edges of its complete graph into at most $n-1$ classes. Since the largest of these classes has size at least 3, Claim 4 and the absence of a universal line together imply that there are points u, v, w, x such that

$$\text{dist}(u, v) = 2, \text{dist}(v, w) = 1, \text{dist}(w, x) = 2 \text{ and } \overline{uv} = \overline{vw} = \overline{wx}$$

or else

$$\text{dist}(v, w) = 1, \text{dist}(w, x) = 2, \text{dist}(u, x) = 1 \text{ and } \overline{vw} = \overline{wx} = \overline{ux}.$$

In both cases, the equality of the three lines implies that

$$\begin{aligned}\text{dist}(u, w) &= \text{dist}(u, x) = \text{dist}(v, w) = \text{dist}(v, x) = 1, \\ \text{dist}(u, v) &= \text{dist}(w, x) = 2.\end{aligned}$$

Since w, x are not twins, there is a point y distinct from both of them and such that $\text{dist}(w, y) \neq \text{dist}(x, y)$; we will complete the proof by showing that $\text{dist}(u, y) \neq \text{dist}(v, y)$.

To do this, assume the contrary: $\text{dist}(u, y) = \text{dist}(v, y)$. Since $y \notin \overline{wx}$ and $\overline{vw} = \overline{wx}$, we have $y \notin \overline{vw}$, and so $\text{dist}(v, y) = \text{dist}(w, y)$. Now $\text{dist}(u, y) \neq \text{dist}(x, y)$, and so $y \in \overline{ux}$; since $y \notin \overline{wx}$, we cannot have $\overline{vw} = \overline{wx} = \overline{ux}$, and so we must have $\overline{uv} = \overline{vw} = \overline{wx}$. In particular, $y \notin \overline{uv}$; since $\text{dist}(u, y) = \text{dist}(v, y)$, we conclude that

$$\text{dist}(u, y) = \text{dist}(v, y) = \text{dist}(w, y) = 2, \quad \text{dist}(x, y) = 1.$$

Since u, v are not twins, there is a point z distinct from both of them and such that $\text{dist}(u, z) \neq \text{dist}(v, z)$; it follows that $\text{dist}(x, z)$ is distinct from one of $\text{dist}(u, z)$, $\text{dist}(v, z)$, and so z belongs to one of the lines $\overline{ux}, \overline{vx}$. But then this line is universal, a contradiction. \square

Claim 9. No critical 1-2 metric space has 5 or 6 points.

Proof. Consider an arbitrary critical 1-2 metric space on n points such that $n = 5$ or $n = 6$ and let u, v, w, x, y be as in Claim 8. We may assume (after a cyclic shift of u, w, v, x if necessary) that

$$\begin{aligned}\text{dist}(u, w) &= \text{dist}(u, x) = \text{dist}(v, w) = \text{dist}(v, x) = 1, \\ \text{dist}(u, v) &= \text{dist}(w, x) = 2, \\ \text{dist}(u, y) &= \text{dist}(w, y) = 1, \quad \text{dist}(v, y) = \text{dist}(x, y) = 2.\end{aligned}$$

Since

$$\overline{ux} \supseteq \{u, v, w, x, y\} \text{ and } \overline{vw} \supseteq \{u, v, w, x, y\},$$

absence of a universal line implies that $n = 6$ and that the sixth point of our space lies outside the lines \overline{ux} and \overline{vw} . Let z denote this sixth point. Since $z \notin \overline{ux}$, $z \notin \overline{vw}$, we have $\text{dist}(u, z) = \text{dist}(x, z)$, $\text{dist}(v, z) = \text{dist}(w, z)$, and so symmetry allows us to distinguish three cases:

- ▷ $\text{dist}(u, z) = \text{dist}(x, z) = 1$, $\text{dist}(v, z) = \text{dist}(w, z) = 1$,
- ▷ $\text{dist}(u, z) = \text{dist}(x, z) = 1$, $\text{dist}(v, z) = \text{dist}(w, z) = 2$,
- ▷ $\text{dist}(u, z) = \text{dist}(x, z) = 2$, $\text{dist}(v, z) = \text{dist}(w, z) = 2$.

Each of these three cases comprises two metric spaces, one with $\text{dist}(y, z) = 1$ and the other with $\text{dist}(y, z) = 2$. Altogether, there are six metric spaces on six points to inspect; each of them has at least six lines. \square

Claim 10. Every metric space on 2, 3, or 4 points has the De Bruijn-Erdős property.

Proof. Consider an arbitrary critical 1-2 metric space on n points with $2 \leq n \leq 4$. If each of its lines has precisely 2 points or if one of its lines has precisely n points, then this space has the De Bruijn-Erdős property; otherwise one of its lines has precisely 3 points and $n = 4$. Let T denote the 3-point line and let w denote the fourth point of the space. If there are distinct x, y in T such that $\overline{wx} = \overline{wy}$, then \overline{xy} is a universal line; else the three lines \overline{wx} with x ranging through T are pairwise distinct 2-point lines. \square

Acknowledgement. I am grateful to the anonymous referee for thoughtfully pointing out several places where my presentation could be, and subsequently was, improved.

References

- [1] *P. Aboulker, A. Bondy, X. Chen, E. Chiniforooshan, P. Miao:* Number of lines in hypergraphs. *Discrete Appl. Math.* *171* (2014), 137–140.
- [2] *X. Chen, V. Chvátal:* Problems related to a De Bruijn-Erdős theorem. *Discrete Appl. Math.* *156* (2008), 2101–2108. [zbl](#) [MR](#)
- [3] *E. Chiniforooshan, V. Chvátal:* A De Bruijn-Erdős theorem and metric spaces. *Discrete Math. Theor. Comput. Sci.* *13* (2011), 67–74. [MR](#)
- [4] *N. G. De Bruijn, P. Erdős:* On a combinatorial problem. *Proc. Akad. Wet. Amsterdam* *51* (1948), 1277–1279. [zbl](#) [MR](#)
- [5] *P. Erdős:* Three point collinearity, Problem 4065. *Am. Math. Mon.* *50* (1943), 65; Solutions in vol. *51* (1944), 169–171.

Author's address: Vašek Chvátal, Department of Computer Science and Software Engineering, Concordia University, Montréal, Canada, e-mail: chvatal@cse.concordia.ca.